

Links of symbolic powers of prime ideals

Hsin-Ju Wang *

Department of Mathematics, National Chung Cheng University
Chiayi 621, Taiwan

Abstract

In this paper, we prove the following. Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Suppose that either R is not regular or R is regular with $d \geq 3$. Let $t \geq 2$ be a positive integer. If $\{\alpha_1, \dots, \alpha_d\}$ is a regular sequence contained in $\mathfrak{m}^t$, then

$$(\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t \subseteq \mathfrak{m}^t.$$

This result gives an affirmative answer to a conjecture raised by Polini and Ulrich.

1 Introduction

Let R be a Noetherian ring. Two proper ideals I and L of height g are said to be *directly linked*, $I \sim L$, if there exists a regular sequence $\mathbf{z} = z_1, \dots, z_g \subseteq I \cap L$ such that $I = (\mathbf{z}) : L$ and $L = (\mathbf{z}) : I$. If R is Cohen-Macaulay and I is an unmixed ideal of height g with R_P Gorenstein for every minimal prime P of I , then one can always produce a link $L \sim I$ by choosing a regular sequence $\mathbf{z} = z_1, \dots, z_g \subset I$ and setting $L = (\mathbf{z}) : I$ ([5]). The linkage class of a proper ideal I is the set of all ideals K obtained by a sequence of links $I = L_0 \sim L_1 \sim \dots \sim L_n = L$, where $n \geq 0$. The algebraic foundations of linkage theory were established in [4] and [5].

In [6], Polini and Ulrich investigated when an ideal is the unique maximal element of its linkage class, in the sense that it contains every ideal of the

*e-mail: hjwang@math.ccu.edu.tw

class. They showed that if $(R, \mathfrak{m})$ is a Gorenstein local ring of dimension $d \geq 2$, with $d \geq 3$ if R is regular, then every ideal in the linkage class of $\mathfrak{m}^t$ is contained in $\mathfrak{m}^t$ provided that $gr_{\mathfrak{m}}(R)$ is Cohen-Macaulay, or R is a complete intersection, or $ecodim R \leq 3$, or $t \leq 3$. They pointed out that $\mathfrak{m}^t$ will be the maximal element of its linkage class if the following conjecture holds.

Conjecture 1: Let $(R, \mathfrak{m})$ be a local Gorenstein ring of dimension $d \geq 2$, with $d \geq 3$ if R is regular, and let $t \geq 1$ be an integer; then every ideal directly linked to $\mathfrak{m}^t$ is contained in $\mathfrak{m}^t$.

In the same paper, they also showed that Conjecture 1 is equivalent to the following question: Is a two-dimensional local Gorenstein ring $(R, \mathfrak{m})$ regular if and only if there exists an integer $t \geq 1$ and a regular sequence $\alpha = \alpha_1, \alpha_2$ contained in $\mathfrak{m}^t$ so that $(\alpha) : \mathfrak{m}^t$ is not contained in $\mathfrak{m}^t$?

In this paper, we settle the conjecture in the affirmative by showing the following:

Theorem 1.1 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$, with $d \geq 3$ if R is regular. Let $t \geq 2$ be a positive integer. If $\{\alpha_1, \dots, \alpha_d\}$ is a regular sequence contained in $\mathfrak{m}^t$, then*

$$(\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t \subseteq \mathfrak{m}^t.$$

Let R be a Noetherian ring. Recall that an ideal I of R of height g is said to be equi-multiple of reduction number 1 if there exists a g generated ideal J contained in I so that $I^2 = JI$. Ideals that are equi-multiple of reduction number 1 enjoy some nice properties: For example, it is shown in [7] that if R is a Cohen-Macaulay local ring and I is an equi-multiple Cohen-Macaulay ideal of reduction number 1, then the associated graded ring $gr_I(R)$ of I is Cohen-Macaulay. Therefore, it is of great interest to search for such classes of ideals. In [1] and [3], Corso, Polini and Vasconcelos proved that the links of prime ideals in Cohen-Macaulay rings are equi-multiple ideals of reduction number 1 under certain assumptions. Moreover, Corso and Polini [2] addressed the following conjecture.

Conjecture 2: Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring and P a prime ideal of height $g \geq 2$. Let $J = (z_1, \dots, z_g)$ be an ideal generated by a regular sequence contained in $P^{(s)}$, where s is a positive integer ≥ 2 . For any

$k = 1, \dots, s$ set $I_k = J : P^{(k)}$. Then

$$I_k^2 = JI_k$$

if one of the following two conditions holds:

(IL_1) R_P is not a regular local ring;

(IL_2) R_P is a regular local ring and two of the z_i 's lie in $P^{(s+1)}$.

With the help of Theorem 1.1 and Theorem 3.2, we are able to give a positive answer to Conjecture 2.

Theorem 1.2 *Let R be a Noetherian ring. Let P be a prime ideal of height $g \geq 2$, $J = (z_1, \dots, z_g)$ an ideal generated by a regular sequence contained in $P^{(k)}$, where $k \geq 2$ is a positive integer. Set $I_k = J : P^{(k)}$. Then $I_k^2 = I_k J$ if one of the following conditions holds:*

(i) R_P is not a regular local ring.

(ii) R_P is a regular local ring and $g \geq 3$.

(iii) R_P is a regular local ring, $g = 2$, and $z_i \in P^{(k+1)}$ for every i .

2 Main Theory

In this section, we study Conjecture 1 stated in section 1. We are able to prove it in a more general form.

Theorem 2.1 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Let I be an ideal of R of height 2 and $\tilde{I} = \sum((\alpha, \beta) : I)$, where the sum is taken over all regular sequences $\{\alpha, \beta\}$ in I . Let t be a positive integer. If $\{\alpha, \beta\}$ is a regular sequence contained in I^t , then*

$$(\alpha, \beta) : I^t \subseteq I^{t-1} \tilde{I}.$$

Remark 2.2 (i) $I \subseteq \tilde{I} \subseteq R$.

(ii) Conjecture 1 follows if we set $d = 2$ and $I = \mathfrak{m}$ in Theorem 2.1 and observe that $\tilde{I} = I = \mathfrak{m}$ if R is not regular.

(iii) One can see from the proof of Theorem 2.1 that the assumption that R is Cohen-Macaulay can be removed if the grade of I is 2.

To show Theorem 2.1, we begin with the following lemmas.

Lemma 2.3 *Let $(R, \mathfrak{m})$ be a Noetherian local ring. Let $\{x_1, \dots, x_{t-1}, y_1, \dots, y_{t-1}\} \subseteq \mathfrak{m}$ such that $\{x_i, y_j\}$ is a regular sequence $\forall i, j$, where $t \geq 2$. Then*

$$\begin{aligned} & \cap_{i=1}^{t-1} (x_1 \cdots x_i, y_1 \cdots y_{t-i}) \\ &= \langle \{x_1 \cdots x_{i-1} y_1 \cdots y_{t-i} \mid 1 \leq i \leq t\} \rangle. \end{aligned}$$

Proof. It is enough to show that

$$\begin{aligned} & \cap_{i=1}^k (x_1 \cdots x_i, y_1 \cdots y_{t-i}) \\ &= (x_1 \cdots x_k) + \langle \{x_1 \cdots x_{i-1} y_1 \cdots y_{t-i} \mid 1 \leq i \leq k\} \rangle \end{aligned}$$

for $1 \leq k \leq t-1$. By induction on k , suppose that the equality holds for k .

Then

$$\begin{aligned} & \cap_{i=1}^{k+1} (x_1 \cdots x_i, y_1 \cdots y_{t-i}) \\ &= ((x_1 \cdots x_k) + \langle \{x_1 \cdots x_{i-1} y_1 \cdots y_{t-i} \mid 1 \leq i \leq k\} \rangle) \cap (x_1 \cdots x_{k+1}, y_1 \cdots y_{t-k-1}) \\ &= (x_1 \cdots x_k) \cap (x_1 \cdots x_{k+1}, y_1 \cdots y_{t-k-1}) + \langle \{x_1 \cdots x_{i-1} y_1 \cdots y_{t-i} \mid 1 \leq i \leq k\} \rangle \\ &= (x_1 \cdots x_{k+1}) + \langle \{x_1 \cdots x_{i-1} y_1 \cdots y_{t-i} \mid 1 \leq i \leq k+1\} \rangle. \end{aligned}$$

□

Lemma 2.4 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Let I be an ideal of R of height 2 and $\tilde{I} = \sum ((\alpha, \beta) : I)$, where the sum is taken over all regular sequences $\{\alpha, \beta\}$ in I . Let $t \geq 1$ and $\{x_1, \dots, x_t, y_1, \dots, y_t\} \subseteq I$ such that $\{x_i, y_j\}$ is a regular sequence $\forall i, j$. Then*

$$(x_1 \cdots x_t, y_1 \cdots y_t) : I^t \subseteq I^{t-1} \tilde{I}.$$

Proof. We may assume that $t \geq 2$. First observe that

$$(x_1 \cdots x_t, y_1 \cdots y_t) : I^t \subseteq \cap_{i=1}^{t-1} (x_1 \cdots x_i, y_1 \cdots y_{t-i}).$$

This is because that

$$\begin{aligned} (x_1 \cdots x_t, y_1 \cdots y_t) : I^t &\subseteq (x_1 \cdots x_t, y_1 \cdots y_t) : x_{i+1} \cdots x_t y_{t-i+1} \cdots y_t \\ &= (x_1 \cdots x_i, y_1 \cdots y_{t-i}) \end{aligned}$$

for $1 \leq i \leq t-1$. By Lemma 2.3, every element in $(\prod_i x_i, \prod_i y_i) : I^t$ is of the form

$$\sum_{i=1}^t a_i x_1 \cdots x_{i-1} y_1 \cdots y_{t-i}.$$

Therefore, to show the lemma, it is enough to show that $a_i \in \tilde{I}$ for every i . For this, let $z \in I$; then

$$\left(\sum_{j=1}^t a_j x_1 \cdots x_{j-1} y_1 \cdots y_{t-j}\right) z x_{i+1} \cdots x_t y_{t-i+2} \cdots y_t \in (x_1 \cdots x_t, y_1 \cdots y_t).$$

Observe that

$$(a_j x_1 \cdots x_{j-1} y_1 \cdots y_{t-j}) z x_{i+1} \cdots x_t y_{t-i+2} \cdots y_t \in (x_1 \cdots x_t, y_1 \cdots y_t)$$

if $j \neq i$. Therefore

$$a_i z x_1 \cdots \hat{x}_i \cdots x_t y_1 \cdots \hat{y}_{t-i+1} \cdots y_t \in (x_1 \cdots x_t, y_1 \cdots y_t).$$

It follows that $a_i z \in (x_i, y_{t-i+1})$, which implies that $a_i \in (x_i, y_{t-i+1}) : I \subseteq \tilde{I}$.
□

Proof of Theorem 2.1: We first prove the following statement: If $y_i \in I \forall i$ and $\{\alpha, y_1 \cdots y_t\}$ is a regular sequence contained in I^t , then

$$(\alpha, y_1 \cdots y_t) : I^t \subseteq I^{t-1} \tilde{I}.$$

Observe that the following set generates I^t :

$$S_t = \{z_1 \cdots z_t \mid z_j \in I, \{y_i, z_j\} \text{ is a regular sequence } \forall i, j\}.$$

Let $a \in (\alpha, y_1 \cdots y_t) : I^t$. Let $z_1 \cdots z_t \in S_t$; then there are elements $u(z), v(z) \in R$ such that

$$a z_1 \cdots z_t = u(z) \alpha + v(z) (y_1 \cdots y_t). \quad (1)$$

Therefore if $z'_1 \cdots z'_t$ is any element in S_t then

$$a z'_1 \cdots z'_t = u(z') \alpha + v(z') (y_1 \cdots y_t). \quad (2)$$

From (1), (2) and the fact that $\{\alpha, y_1 \cdots y_t\}$ is a regular sequence, we see that $u(z) z'_1 \cdots z'_t \in (z_1 \cdots z_t, y_1 \cdots y_t)$. Since $z'_1 \cdots z'_t$ is an arbitrary element in S_t ,

$$u(z) \in (z_1 \cdots z_t, y_1 \cdots y_t) : I^t \subseteq I^{t-1} \tilde{I}$$

by Lemma 2.4. Since $z_1 \cdots z_t$ is an arbitrary element in S_t and $\alpha \in I^t$, there are elements $u, v \in R$ with $u \in I^{t-1} \tilde{I}$ such that $a \alpha = u \alpha + v (y_1 \cdots y_t)$, which

implies that $a - u \in (y_1 \cdots y_t)$. It follows that $a \in I^{t-1}\tilde{I}$ and this completes the proof of the statement.

In general, if $\{\alpha, \beta\}$ is a regular sequence contained in I^t , then we can use the same argument the above to conclude that every element in $(\alpha, \beta) : I^t$ is in $I^{t-1}\tilde{I}$.

Corollary 2.5 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Suppose that either R is not regular or R is regular with $d \geq 3$. Let $t \geq 2$ be a positive integer. If $\{\alpha_1, \dots, \alpha_d\}$ is a regular sequence contained in $\mathfrak{m}^t$, then*

$$(\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t \subseteq \mathfrak{m}^t.$$

Proof. We prove by induction on d . The case $d = 2$ follows from Theorem 2.1. Assume that $d \geq 3$. Let $a \in (\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t$. Since $R/(\alpha_d)$ is Cohen-Macaulay and is not regular, $a \in \mathfrak{m}^t \pmod{(\alpha_d)}$ by induction. It follows that $a \in \mathfrak{m}^t$ as $\alpha_d \in \mathfrak{m}^t$. $\square$

The following result will be used in section 3.

Corollary 2.6 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Suppose that either R is not regular or R is regular with $d \geq 3$. Let $t \geq 2$ be a positive integer and $\{\alpha_1, \dots, \alpha_d\}$ be a regular sequence contained in $\mathfrak{m}^t$. Then for every $a \in (\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t$ and every element $f \in \mathfrak{m}^t$, if $af = \sum_{i=1}^d u_i \alpha_i$ then $u_i \in \mathfrak{m}^t$ for every i .*

Corollary 2.6 follows from the next lemma.

Lemma 2.7 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Let $t \geq 2$ be a positive integer. Assume that for any regular sequence $\{\alpha_1, \dots, \alpha_d\}$ contained in $\mathfrak{m}^t$, $(\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t \subseteq \mathfrak{m}^t$. Then for every $a \in (\alpha_1, \dots, \alpha_d) : \mathfrak{m}^t$ and every element $f \in \mathfrak{m}^t$, if $af = \sum_{i=1}^d u_i \alpha_i$ then $u_i \in \mathfrak{m}^t$ for every i .*

Proof. It is easy to see that the following set generates the ideal $\mathfrak{m}^t$:

$$S_t = \{z_1 \cdots z_t \mid z_j \in \mathfrak{m}, \{\alpha_1, \dots, \hat{\alpha}_i, \dots, \alpha_d, z_j\} \text{ is a regular sequence } \forall i, j\}.$$

Let $z_1 \cdots z_t \in S_t$; then there are $u_i(z) \in R$ such that

$$az_1 \cdots z_t = \sum_{i=1}^d u_i(z) \alpha_i.$$

From the proof of Theorem 2.1, we see that $u_i(z) \in (\alpha_1, \dots, \hat{\alpha}_i, \dots, \alpha_d, z_1 \cdots z_t) : \mathfrak{m}^t$ for every i . Therefore $u_i(z) \in \mathfrak{m}^t$ by assumption. This shows that if $f \in \mathfrak{m}^t$ then af can be expressed as $\sum_{i=1}^d u_i \alpha_i$ for some $u_i \in \mathfrak{m}^t$ for every i . If $af = \sum_{i=1}^d u'_i \alpha_i$ for some $u'_i \in R$, then $u_i - u'_i \in (\alpha_1, \dots, \hat{\alpha}_i, \dots, \alpha_d) \subseteq \mathfrak{m}^t$, it follows that $u'_i \in \mathfrak{m}^t$ for every i . $\square$

3 Ideals of reduction number 1

The goal of this section is to study the following conjecture:

Conjecture: Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring and P a prime ideal of height $g \geq 2$. Let $J = (z_1, \dots, z_g)$ be an ideal generated by a regular sequence contained in $P^{(s)}$, where s is a positive integer ≥ 2 . For any $k = 1, \dots, s$ set $I_k = J : P^{(k)}$. Then

$$I_k^2 = JI_k$$

if one of the following two conditions holds:

(IL₁) R_P is not a regular local ring;

(IL₂) R_P is a regular local ring and two of the z_i 's lie in $P^{(s+1)}$.

There are some partial answers to this conjecture in [2] and [6]. The following result solves the conjecture completely.

Theorem 3.1 *Let R be a Noetherian ring. Let P be a prime ideal of height $g \geq 2$, $J = (z_1, \dots, z_g)$ an ideal generated by a regular sequence contained in $P^{(k)}$, where $k \geq 2$ is a positive integer. Set $I_k = J : P^{(k)}$. Then $I_k^2 = I_k J$ if one of the following conditions holds:*

(i) R_P is not a regular local ring.

(ii) R_P is a regular local ring and $g \geq 3$.

(iii) R_P is a regular local ring, $g = 2$, and $z_i \in P^{(k+1)}$ for every i .

To show Theorem 3.1, we need the following result.

Theorem 3.2 *Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension $d \geq 2$. Let $k \geq 2$ be a positive integer. If $\mathbf{z} = z_1, \dots, z_d$ is a regular sequence contained in $\mathfrak{m}^{k+1}$, then*

$$(\mathbf{z}) : \mathfrak{m}^k \subseteq (\mathfrak{m}^{2k+1} : \mathfrak{m}^k)$$

if $\text{depth } gr_{\mathfrak{m}}(R) \geq 2$.

Proof. It is not difficult to see that the following set generates the ideal $\mathfrak{m}^k$.

$$S_k = \left\{ \prod_{i=1}^k y_i \mid y_i \in \mathfrak{m}, \{y_i^*, y_j^*\} \text{ is a regular sequence in } gr_{\mathfrak{m}}(R) \right\}.$$

Let $a \in (\mathbf{z}) : \mathfrak{m}^k$ and $f = \prod_{i=1}^k y_i \in S_k$; then

$$af = \sum_{i=1}^d u_i z_i \tag{3}$$

for some $u_i \in R$. To show the theorem it is enough to show that $u_i \in \mathfrak{m}^k$ as $z_i \in \mathfrak{m}^{k+1}$ for every i . Without loss of generality, we only prove that $u_1 \in \mathfrak{m}^k$ as an example.

Choose $y \in \mathfrak{m}$ so that $\{y_j^*, y^*\}$ form a regular sequence in $gr_{\mathfrak{m}}(R)$ for every j . Let $f_j = yf/y_j$; then $f \in S_k$, therefore there are $u_{ij} \in R$ such that

$$af_j = \sum_{i=1}^d u_{ij} z_i. \tag{4}$$

From (3) and (4), we obtain that

$$(u_1 y - u_{1j} y_j) z_1 \in (z_2, \dots, z_d).$$

Moreover, since $\{z_1, \dots, z_d\}$ is a regular sequence and $z_i \in \mathfrak{m}^{k+1}$ by assumption,

$$(u_1 y - u_{1j} y_j) \in \mathfrak{m}^{k+1}.$$

It follows that

$$u_1 \in (y_j) + \mathfrak{m}^k$$

for every j by the choice of y . Now one can conclude that $u_1 \in (y_l \cdots y_k) + \mathfrak{m}^k$ by the facts that $\{y_l^*, y_j^*\}$ is a regular sequence in $gr_{\mathfrak{m}}(R)$ if $j > l$ for $l = k-1, k-2, \dots$, in particular, $u_1 \in \mathfrak{m}^k$.

□

Corollary 3.3 *Let $(R, \mathfrak{m})$ be a 2-dimensional regular local ring. Let $k \geq 2$ and $\mathbf{z} = z_1, z_2$ be regular sequence contained in $\mathfrak{m}^{k+1}$. Then the following hold.*

$$(i) \ (\mathbf{z}) : \mathfrak{m}^k \subseteq \mathfrak{m}^k;$$

$$(ii) \text{ For every element } a \in (\mathbf{z}) : \mathfrak{m}^k \text{ and every element } f \in \mathfrak{m}^k, \text{ if } af = \sum_{i=1}^2 u_i z_i \text{ then } u_i \in \mathfrak{m}^k.$$

Proof. (i) follows from the fact that $gr_{\mathfrak{m}}(R)$ is Cohen-Macaulay and $\mathfrak{m}^{2k+1} : \mathfrak{m}^k = \mathfrak{m}^{k+1} \subseteq \mathfrak{m}^k$. (ii) follows from the proof of Theorem 3.2. □

Before proving Theorem 3.1, we need one more lemma.

Lemma 3.4 *Let R be a Noetherian ring. Let H be an ideal of height $g \geq 2$, $J = (z_1, \dots, z_g)$ an ideal generated by a regular sequence contained in H . Set $I = J : H$. If $I = (a_1, \dots, a_t)$, $a_i H \subseteq JH$ for every i , and $I^2 \subseteq J$, then $I^2 = IJ$.*

Proof. Since $I^2 \subseteq J$, there are $u_{ij}^{(k)} \in R$ such that $a_i a_j = \sum_{k=1}^g u_{ij}^{(k)} z_k$. To complete the proof it suffices to show that $u_{ij}^{(k)} \in I$, i.e., $u_{ij}^{(k)} H \subseteq J$ for every k . For this let $b \in H$; then by assumption,

$$\begin{aligned} \sum_{k=1}^g (u_{ij}^{(k)} b) z_k &= a_i a_j b \\ &\in a_i JH \\ &\subseteq J^2 H \\ &\subseteq (z_1, \dots, z_g)^2. \end{aligned}$$

Since $z_1, \dots, z_g$ is a regular sequence (although not necessarily permutable), we see that $u_{ij}^{(k)} b \in (z_1, \dots, z_g)$ for every k . □

Proof of Theorem 3.1: Let $I = I_k = J : P^{(k)}$. Then $I \subseteq P^{(k)}$ by

Corollary 2.5 and Corollary 3.3(i), it follows that $I^2 \subseteq IP^{(k)} \subseteq J$. To complete the proof, by Lemma 3.4, we need only to prove that if $a \in I$, $f \in P^{(k)}$ and $af = \sum_{i=1}^g u_i z_i$, then $u_i \in P^{(k)}$ for every i . However this follows by Corollary 2.6 and Corollary 3.3(ii).

4 The case $t = 1$

We study the links of unmixed radical ideals in a Noetherian ring in this section. The main result Theorem 4.2 will generalize [1, Theorem 2.3] and [3, Theorem 2.1].

Throughout, let R be a Noetherian ring of depth $d \geq g$, H an unmixed radical ideal of height g , and let $\mathbf{z} = z_1, \dots, z_g \subseteq H$ be a regular sequence. Set $J = (\mathbf{z})$ and $I = J : H$. Suppose that for every minimal prime P of H , one of the following two conditions holds:

- (L_1) R_P is not a regular local ring.
- (L_2) R_P is a regular local ring of dimension at least 2 and two elements in the sequence $\mathbf{z} = z_1, \dots, z_g$ lie in the symbolic square $P^{(2)}$.

An easy observation is the following.

Lemma 4.1 $I \subseteq H$ and $I^2 \subseteq J$.

Proof. To show $I \subseteq H$, it suffices to show that $I \subseteq P$ for every minimal prime P of H . However, if I is not a subset of P then $P_P = J_P$, which contradicts to L_1 or L_2 . Thus $I \subseteq H$ is fulfilled. Moreover, $I^2 \subseteq IH \subseteq J$. $\square$

Theorem 4.2 *Let R be a Noetherian ring of depth $d \geq g$, H an unmixed radical ideal of height g , and $\mathbf{z} = z_1, \dots, z_g \subseteq H$ a regular sequence. Set $J = (\mathbf{z})$ and $I = J : H$. Suppose that for every minimal prime P of H , either L_1 or L_2 holds. Then $I^2 = JI$.*

Proof. Write $H = P_1 \cap \dots \cap P_l$. For $1 \leq j \leq l$, let

$$\begin{aligned} T_j &= \cup_i \text{Ass}(R/(z_1, \dots, \hat{z}_i, \dots, z_g)) \cap \text{Spec}(R_{P_j}) \\ &= \cup_i \text{Min}(R/(z_1, \dots, \hat{z}_i, \dots, z_g)) \cap \text{Spec}(R_{P_j}). \end{aligned}$$

Since I is not a subset of every prime in $\cup_j T_j$, we can choose a generating set $\{a_1, \dots, a_t\}$ of I so that $a_i \notin \cup_j T_j$ for every i .

Now, to finish the proof, it is enough to show that $a_i H \subseteq JH$ for every i by

Lemma 3.4. For this, let $b \in H$ and a be any a_i ; then there are elements $u_j \in R$ such that

$$ab = \sum_{j=1}^g u_j z_j. \quad (5)$$

To show $u_j \in H$, we can prove it locally. Without loss of generality, we will prove that $u_1 \in PR_P$ for every minimal prime P of H .

Let b' be an arbitrary element in H ; then there are elements $u'_j \in R$ such that

$$ab' = \sum_{j=1}^g u'_j z_j. \quad (6)$$

From (5) and (6), we obtain that

$$a(b'u_1 - bu'_1) \in (z_2, \dots, z_g).$$

Let P be any minimal prime of H ; then $\{a, z_2, \dots, z_g\}$ is a regular sequence in R_P . Since b' is arbitrary, $u_1 P_P = u_1 H_P \subseteq (b, z_2, \dots, z_g)R_P$. From the fact that R_P is not regular or at least one element in $\{z_2, \dots, z_g\}$ is in $P^{(2)}$, we conclude that $u_1 \in PR_P$. $\square$

Acknowledgements

The author would like to thank C. Huneke for helpful suggestion which improved the exposition of Theorem 2.1.

References

- [1] A. Corso and C. Polini, *Links of prime ideals and their Rees algebras*, J. Algebra **178** (1995), 224–238.
- [2] A. Corso and C. Polini, *Reduction number of links of irreducible varieties*, J. Pure and Applied Algebra **121** (1997), 29–43.
- [3] A. Corso, C. Polini and W.V. Vasconcelos, *Links of prime ideals*, Math. Proc. Cambridge Philos. Soc. **115** (1994), 431–436.
- [4] C. Huneke, and B. Ulrich, *The structure of linkage*, Ann. Math. **126** (1987), 277–334.

- [5] C. Peskine and L. Szpiro, *Liaison des variétés algébriques*, Invent. Math. **26** (1974), 271–302.
- [6] C. Polini and B. Ulrich, *Linkage and reduction numbers*, Math. Annalen **310** (1998), 631–651.
- [7] W.V. Vasconcelos, *Hilbert functions, analytic spread and Koszul homology*, in Commutative Algebra: Syzygies, Multiplicities, and Birational Algebra, W. Heinzer, C. Huneke, J. Sally (eds.), Contemp. Math. **159**, Amer. Math. Soc., Providence, (1994), 401–422.